

## OPIOID PRODUCTS

GENERIC	BRAND	DEA SCHEDULE	DOSAGE FORMS	ADDITIONAL COMMENTS
<b>MILD TO MODERATELY SEVERE PAIN</b>				
Codeine sulfate		Schedule II	<b>Tab:</b> 15mg, 30mg, 60mg; <b>Sol:</b> 30mg/5mL	<ul style="list-style-type: none"> <li>• <b>BW:</b> Death related to ultra-rapid metabolism of codeine to morphine</li> <li>• Pregnancy category C, caution in nursing</li> </ul>
Codeine phosphate/Acetaminophen		Schedule V	<b>Sol:</b> 12mg-120mg/5mL	<ul style="list-style-type: none"> <li>• <b>BW:</b> Hepatotoxicity</li> <li>• Pregnancy category C, not for use in nursing</li> </ul>
		Schedule III	<b>Tab:</b> 15mg-300mg	
	Tylenol #3	Schedule III	<b>Tab:</b> 30mg-300mg	
	Tylenol #4	Schedule III	<b>Tab:</b> 60mg-300mg	
<b>MODERATE TO MODERATELY SEVERE PAIN</b>				
Dihydrocodeine bitartrate/Aspirin/Caffeine	Synalgos-DC	Schedule III	<b>Cap:</b> 16mg-356.4mg-30mg	<ul style="list-style-type: none"> <li>• <b>BW:</b> Death related to ultra-rapid metabolism of codeine to morphine</li> <li>• Safety not known in pregnancy, not for use in nursing</li> </ul>
Hydrocodone bitartrate/Acetaminophen	Norco	Schedule III	<b>Tab:</b> 5mg-325mg, 7.5mg-325mg, 10mg-325mg	<ul style="list-style-type: none"> <li>• <b>BW:</b> Hepatotoxicity</li> <li>• Pregnancy category C, not for use in nursing</li> </ul>
	Vicodin		<b>Tab:</b> 5mg-300mg	
	Vicodin ES		<b>Tab:</b> 7.5mg-300mg	
	Vicodin HP		<b>Tab:</b> 10mg-300mg	
Hydrocodone bitartrate/Ibuprofen	Reprexain	Schedule III	<b>Tab:</b> 2.5mg-200mg, 5mg-200mg, 10mg-200mg	<ul style="list-style-type: none"> <li>• Pregnancy category C, not for use in nursing</li> </ul>
	Vicoprofen		<b>Tab:</b> 7.5-200mg	
Oxycodone HCl/Acetaminophen	Percocet	Schedule II	<b>Tab:</b> 2.5mg-325mg, 5mg-325mg, 7.5mg-325mg, 10mg-325mg	<ul style="list-style-type: none"> <li>• <b>BW:</b> Hepatotoxicity</li> <li>• Pregnancy category C, not for use in nursing</li> </ul>
	Roxicet		<b>Tab:</b> 5mg-325mg; <b>Sol:</b> 5mg-325mg/5mL	
Oxycodone HCl/Aspirin	Percodan	Schedule II	<b>Tab:</b> 4.8355mg-325mg	<ul style="list-style-type: none"> <li>• Pregnancy category B (oxycodone) and D (aspirin), not for use in nursing</li> </ul>

(Continued)

GENERIC	BRAND	DEA SCHEDULE	DOSAGE FORMS	ADDITIONAL COMMENTS
<b>MODERATE TO SEVERE OR CHRONIC PAIN</b>				
<b>Buprenorphine</b>	Butrans	Schedule III	<b>Patch:</b> 5mcg/hr, 10mcg/hr, 15mcg/hr, 20mcg/hr	<ul style="list-style-type: none"> <li>• <b>BW:</b> Abuse potential, life-threatening respiratory depression, and accidental exposure</li> <li>• Pregnancy category C, not for use in nursing</li> </ul>
<b>Fentanyl</b>	Duragesic	Schedule II	<b>Patch:</b> 12mcg/hr, 25mcg/hr, 50mcg/hr, 75mcg/hr, 100mcg/hr	<ul style="list-style-type: none"> <li>• <b>BW:</b> Abuse potential, life-threatening respiratory depression, and accidental exposure</li> <li>• Pregnancy category C, not for use in nursing</li> </ul>
<b>Hydromorphone HCl</b>	Dilaudid, Dilaudid-HP	Schedule II	<b>Tab:</b> 2mg, 4mg, 8mg; <b>Sol:</b> 1mg/mL; <b>Inj:</b> 1mg/mL, 2mg/mL, 4mg/mL; <b>HP: Inj:</b> 10mg/mL [1mL, 5mL, 50mL], 250mg (sterile, lyophilized powder)	<ul style="list-style-type: none"> <li>• <b>BW:</b> Risk of respiratory depression, abuse, and medication errors. Dilaudid-HP injection is for use in opioid-tolerant patients only</li> <li>• Pregnancy category C, not for use in nursing</li> </ul>
	Exalgo		<b>Tab, ER:</b> 8mg, 12mg, 16mg, 32mg	
<b>Levorphanol tartrate</b>		Schedule II	<b>Tab:</b> 2mg	<ul style="list-style-type: none"> <li>• Pregnancy category C, not for use in nursing</li> </ul>
<b>Meperidine HCl</b>		Schedule II	<b>Sol:</b> 50mg/5mL	<ul style="list-style-type: none"> <li>• Pregnancy category C, not for use in nursing</li> </ul>
	Demerol		<b>Tab:</b> 50mg, 100mg; <b>Inj:</b> 25mg/mL, 50mg/mL, 75mg/mL, 100mg/mL	
<b>Methadone HCl</b>		Schedule II	<b>Inj:</b> 10mg/mL	<ul style="list-style-type: none"> <li>• <b>BW:</b> Life-threatening QT prolongation</li> <li>• Pregnancy category C, not for use in nursing</li> </ul>
			<b>Sol:</b> 5mg/5mL, 10mg/5mL	<ul style="list-style-type: none"> <li>• <b>BW:</b> Abuse potential, life-threatening respiratory depression, life-threatening QT prolongation, accidental exposure, and treatment for opioid addiction</li> <li>• Pregnancy category C, caution in nursing</li> </ul>
	Dolophine		<b>Tab:</b> 5mg, 10mg	
	Methadose		<b>Sol, Concentrated:</b> 10mg/mL; <b>Tab:</b> 5mg, 10mg; <b>Tab, Dispersible:</b> 40mg	
<b>Morphine sulfate/Naltrexone HCl</b>	Embeda	Schedule II	<b>Cap, ER:</b> 20mg-0.8mg, 30mg-1.2mg, 50mg-2mg, 60mg-2.4mg, 80mg-3.2mg, 100mg-4mg	<ul style="list-style-type: none"> <li>• <b>BW:</b> Abuse potential, life-threatening respiratory depression, accidental exposure, and interaction with alcohol</li> <li>• Pregnancy category C, not for use in nursing</li> </ul>
<b>Morphine sulfate</b>		Schedule II	<b>Tab:</b> 15mg, 30mg; <b>Sol:</b> 10mg/5mL, 20mg/5mL, 100mg/5mL; <b>Inj:</b> 2mg/mL, 4mg/mL, 8mg/mL, 10mg/mL, 15mg/mL	<ul style="list-style-type: none"> <li>• <b>BW:</b> (Sol) Risk of medication errors. The 100mg/5mL is indicated for opioid-tolerant patients only</li> <li>• Pregnancy category C, not for use in nursing</li> </ul>
	Astramorph/PF		<b>Inj:</b> 0.5mg/mL, 1mg/mL	<ul style="list-style-type: none"> <li>• Pregnancy category C, safety not known in nursing</li> </ul>

(Continued)

GENERIC	BRAND	DEA SCHEDULE	DOSAGE FORMS	ADDITIONAL COMMENTS
<b>MODERATE TO SEVERE OR CHRONIC PAIN</b> <i>(Continued)</i>				
<b>Morphine sulfate</b> <i>(Continued)</i>	Avinza	Schedule II	<b>Cap, ER:</b> 30mg, 45mg, 60mg, 75mg, 90mg, 120mg	<ul style="list-style-type: none"> <li>• <b>BW:</b> Abuse potential, life-threatening respiratory depression, accidental exposure, and interaction with alcohol</li> <li>• Pregnancy category C, not for use in nursing</li> </ul>
	Duramorph		<b>Inj:</b> 0.5mg/mL, 1mg/mL	<ul style="list-style-type: none"> <li>• <b>BW:</b> Risk of severe adverse reactions with epidural or intrathecal route; observe patients in an equipped and staffed environment for at least 24 hrs after initial dose</li> <li>• Pregnancy category C, safety not known in nursing</li> </ul>
	Infumorph		<b>Inj:</b> 0.5mg/mL (5mg/10mL), 1mg/mL (10mg/10mL), 10mg/mL (200mg/20mL), 25mg/mL (500mg/20mL)	<ul style="list-style-type: none"> <li>• <b>BW:</b> Not recommended for single-dose IV, IM, or SQ administration. Risk of severe adverse reactions; observe patients in an equipped and staffed environment for at least 24 hrs after initial (single) test dose, and as appropriate, for the 1st several days after catheter implantation</li> <li>• Pregnancy category C, safety not known in nursing</li> </ul>
	Kadian		<b>Cap, ER:</b> 10mg, 20mg, 30mg, 40mg, 50mg, 60mg, 70mg, 80mg, 100mg, 130mg, 150mg, 200mg	<ul style="list-style-type: none"> <li>• <b>BW:</b> Abuse potential, life-threatening respiratory depression, and accidental exposure</li> <li>• Pregnancy category C, not for use in nursing</li> </ul>
	MS Contin		<b>Tab, ER:</b> 15mg, 30mg, 60mg, 100mg, 200mg	<ul style="list-style-type: none"> <li>• <b>BW:</b> Abuse potential, life-threatening respiratory depression, and accidental exposure</li> <li>• Pregnancy category C, not for use in nursing</li> </ul>
<b>Oxycodone HCl</b>		Schedule II	<b>Cap:</b> 5mg; <b>Sol:</b> 1mg/mL, 20mg/mL	<ul style="list-style-type: none"> <li>• <b>BW:</b> (Sol) Risk of medication errors. The 100mg/5mL is indicated for opioid-tolerant patients only</li> <li>• Pregnancy category B, not for use in nursing</li> </ul>
	Oxecta		<b>Tab:</b> 5mg, 7.5mg	<ul style="list-style-type: none"> <li>• <b>BW:</b> (OxyContin) Abuse potential, life-threatening respiratory depression, and accidental exposure</li> <li>• Pregnancy category B, not for use in nursing</li> </ul>
	Roxicodone		<b>Tab:</b> 5mg, 15mg, 30mg	
	OxyContin		<b>Tab, ER:</b> 10mg, 15mg, 20mg, 30mg, 40mg, 60mg, 80mg	
<b>Oxymorphone HCl</b>	Opana	Schedule II	<b>Inj:</b> 1mg/mL; <b>Tab:</b> 5mg, 10mg	<ul style="list-style-type: none"> <li>• Pregnancy category C, caution in nursing</li> </ul>
	Opana ER		<b>Tab, ER:</b> 5mg, 7.5mg, 10mg, 15mg, 20mg, 30mg, 40mg	<ul style="list-style-type: none"> <li>• <b>BW:</b> Abuse potential, life-threatening respiratory depression, accidental exposure, and interaction with alcohol</li> <li>• Pregnancy category C, caution in nursing</li> </ul>
<b>Tapentadol</b>	Nucynta	Schedule II	<b>Tab:</b> 50mg, 75mg, 100mg	<ul style="list-style-type: none"> <li>• <b>BW:</b> (Nucynta ER) Abuse potential, life-threatening respiratory depression, accidental exposure, and interaction with alcohol</li> <li>• Pregnancy category C, not for use in nursing</li> </ul>
	Nucynta ER		<b>Tab, ER:</b> 50mg, 100mg, 150mg, 200mg, 250mg	

*(Continued)*

GENERIC	BRAND	DEA SCHEDULE	DOSAGE FORMS	ADDITIONAL COMMENTS
<b>BREAKTHROUGH PAIN</b>				
<b>Fentanyl</b>	Abstral	Schedule II	<b>Tab, SL:</b> 100mcg, 200mcg, 300mcg, 400mcg, 600mcg, 800mcg	<ul style="list-style-type: none"> <li>• <b>BW:</b> Risk of respiratory depression, medication errors, and abuse potential</li> <li>• Pregnancy category C, not for use in nursing</li> </ul>
	Fentora		<b>Tab, Buccal:</b> 100mcg, 200mcg, 400mcg, 600mcg, 800mcg	
	Lazanda		<b>Nasal:</b> 100mcg/spr, 400mcg/spr	
	Subsys		<b>Spr, SL:</b> 100mcg/spr, 200mcg/spr, 400mcg/spr, 600mcg/spr, 800mcg/spr, 1200mcg/spr, 1600mcg/spr	
<b>Fentanyl citrate</b>	Actiq	Schedule II	<b>Loz:</b> 200mcg, 400mcg, 600mcg, 800mcg, 1200mcg, 1600mcg	<ul style="list-style-type: none"> <li>• <b>BW:</b> Risk of respiratory depression, medication errors, and abuse potential</li> <li>• Pregnancy category C, not for use in nursing</li> </ul>
Refer to full FDA-approved labeling for more detailed dosing.				
<b>Abbreviations:</b> BW = boxed warning; ER = extended-release; SL = sublingual.				