

ACNE MANAGEMENT

GENERIC (BRAND)	HOW SUPPLIED	DOSAGE
SYSTEMIC AGENTS		
Isotretinoin (Absorica)	Cap: 10mg, 20mg, 25mg, 30mg, 35mg, 40mg	Adults & Peds ≥12 yrs: 0.5-1mg/kg/day given in 2 divided doses for 15-20 wks. Max: 2mg/kg/day.
Isotretinoin (Amnesteem, Claravis)	Cap: (Amnesteem) 10mg, 20mg, 40mg; (Claravis) 10mg, 20mg, 30mg, 40mg	Adults & Peds ≥12 yrs: 0.5-1mg/kg/day given in 2 divided doses for 15-20 wks. Max: 2mg/kg/day.
Minocycline HCl (Solodyn)	Tab, Extended-Release: 55mg, 65mg, 80mg, 105mg, 115mg	Adults & Peds ≥12 yrs: 1mg/kg qd for 12 wks.
TOPICAL AGENTS		
Adapalene (Differin)	Cre: 0.1%	Adults & Peds ≥12 yrs: Apply a thin layer to affected area qpm.
	Gel: 0.1%, 0.3%	Adults & Peds ≥12 yrs: Apply a thin layer to affected area qhs after washing.
	Lot: 0.1%	Adults & Peds ≥12 yrs: Apply a thin layer to cover the entire face and other affected area qd after washing.
Azelaic acid (Azelex)	Cre: 20%	Adults & Peds ≥12 yrs: Wash and dry skin. Massage gently but thoroughly into affected area bid (am and pm).
Benzoyl peroxide (BenzEfoam, BenzEfoam Ultra)	(BenzEfoam) Foam: 5.3%	Adults & Peds ≥12 yrs: Rub into affected area qd or ud until completely absorbed. Facial Acne: Use a dollop the size of a marble. Back/Chest Acne: Use a dollop the size of a whole walnut.
	(BenzEfoam Ultra) Foam: 9.8%	Adults & Peds ≥12 yrs: Rub into affected area qd, then bid-tid prn or ud until completely absorbed. Rinse off after 2 min.
Benzoyl peroxide (NeoBenz Micro SD, NeoBenz Micro Wash)	(SD) Cre: 5.5%	Adults & Peds ≥12 yrs: Apply to affected area qd-bid.
	(Wash) Sol: 7%	
Benzoyl peroxide	Gel: 2.5%, 5%, 10%	Adults & Peds ≥12 yrs: (2.5%) Initial: Apply to affected area qd for 1 week, then bid. (5%, 10%) May be initiated in patients demonstrating accommodation to 2.5% gel.
Clindamycin phosphate (Evoclin)	Foam: 1%	Adults & Peds ≥12 yrs: Apply to affected area qd.
Clindamycin phosphate (Cleocin T)	Gel: 1%	Adults & Peds ≥12 yrs: Apply thin layer to affected area bid.
	Lot: 1%	
	Sol: 1%	
Clindamycin phosphate (Clindagel)	Gel: 1%	Adults & Peds ≥12 yrs: Apply thin layer to affected area qd.
Dapsone (Aczone)	Gel: 5%	Adults & Peds ≥12 yrs: Apply pea-sized amount to affected area bid.
Erythromycin (Akne-Mycin)	Oint: 2%	Adults: Apply to affected area bid (am and pm).
Erythromycin	Gel: 2%	Adults: Wash and dry skin. Apply a thin layer to affected area qd-bid. Do not rub in; spread lightly.
	Pledgets: 2%	Adults: Wash and dry skin. Rub pledget or apply sol to affected area bid (am and pm).
	Sol: 2%	
Gentamicin sulfate	Cre: 0.1%	Adults & Peds >1 yr: Apply small amount to affected area tid-qid.
	Oint: 0.1%	

(Continued)

GENERIC (BRAND)	HOW SUPPLIED	DOSAGE
TOPICAL AGENTS <i>(Continued)</i>		
Tazarotene (Fabior)	Foam: 0.1%	Adults & Peds ≥12 yrs: Wash and dry skin. Apply thin layer to affected area of face and/or upper trunk qpm. Massage into skin.
Tazarotene (Tazorac)	Cre: 0.1% Gel: 0.1%	Adults & Peds ≥12 yrs: Wash and dry skin. Apply thin layer to affected area qpm.
Tretinoin (Atralin)	Gel: 0.05%	Adults & Peds ≥10 yrs: Apply thin layer to affected area qhs.
Tretinoin (Avita)	Cre: 0.025% Gel: 0.025%	Adults: Apply thin layer to affected area qpm.
Tretinoin (Retin-A)	Cre: 0.025%, 0.05%, 0.1% Gel: 0.01%, 0.025%	Adults: Apply thin layer to affected area qhs.
Tretinoin (Retin-A Micro)	Gel: 0.04%, 0.08%, 0.1%	Adults & Peds ≥12 yrs: Apply thin layer to affected area qpm.
Tretinoin (Tretin-X)	Cre: 0.025%, 0.0375%, 0.05%, 0.075%, 0.1%	Adults & Peds ≥12 yrs: Apply thin layer to affected area qhs.
COMBINATION PRODUCTS, TOPICAL		
Adapalene-Benzoyl peroxide (Epiduo)	Gel: 0.1%-2.5%	Adults & Peds ≥9 yrs: Apply pea-sized amount to affected area of face and/or trunk qd after washing.
Benzoyl peroxide-Clindamycin phosphate (Acanya)	Gel: 2.5%-1.2%	Adults & Peds ≥12 yrs: Apply pea-sized amount to face qd.
Benzoyl peroxide-Clindamycin phosphate (BenzaClin)	Gel: 5%-1%	Adults & Peds ≥12 yrs: Wash and dry skin. Apply to affected area bid or ud.
Benzoyl peroxide-Clindamycin phosphate (Duac)	Gel: 5%-1.2%	Adults & Peds ≥12 yrs: Wash and dry skin. Apply thin layer to face qpm or ud.
Benzoyl peroxide-Erythromycin (Benzamycin)	Gel: 5%-3%	Adults & Peds ≥12 yrs: Wash and dry skin. Apply to affected area bid or ud.
Clindamycin phosphate-Tretinoin (Veltin)	Gel: 1.2%-0.025%	Adults & Peds ≥12 yrs: Apply pea-sized amount lightly covering the entire affected area qpm.
Clindamycin phosphate-Tretinoin (Ziana)	Gel: 1.2%-0.025%	Adults & Peds ≥12 yrs: Apply pea-sized amount onto 1 fingertip, dot onto chin, cheeks, nose, and forehead, then gently rub over entire face qhs.
Hydrocortisone acetate-Iodoquinol (Alcortin-A)	Gel: 2%-1%	Adults & Peds ≥12 yrs: Apply to affected area tid-qid or ud.
Sodium sulfacetamide (Klaron)	Lot: 10%	Adults & Peds ≥12 yrs: Apply thin layer to affected area bid.
Sodium sulfacetamide-Sulfur (Avar Cleanser, Avar Cleansing Pads, Avar LS Cleanser, Avar LS Cleansing Pads, Avar-e Cream, Avar-e Green Cream, Avar-e LS Cream)	Pads: (Avar Cleansing Pads) 9.5%-5%; (Avar LS Cleansing Pads) 10%-2%	Adults & Peds ≥12 yrs: Wash affected area qd-bid or ud.
	Cre: (Avar-e Cream, Avar-e Green Cream) 10%-5%; (Avar-e LS Cream) 10%-2%	Adults & Peds ≥12 yrs: Cleanse affected area. Apply a thin layer to affected area qd-tid or ud. Massage into skin.
	Sol: (Avar Cleanser) 10%-5%; (Avar LS Cleanser) 10%-2%	Adults & Peds ≥12 yrs: Wash affected area qd-bid or ud.
Sodium sulfacetamide-Sulfur (Sumadan)	Wash: 9%-4.5%	Adults & Peds ≥12 yrs: Apply to affected area qd-bid or ud.

GENERIC (BRAND)	HOW SUPPLIED	DOSAGE
COMBINATION PRODUCTS, TOPICAL <i>(Continued)</i>		
Sodium sulfacetamide-Sulfur	Lot: 10%-5%	Adults & Peds ≥12 yrs: Apply thin layer to affected area with light massaging qd-tid or ud.
Refer to full FDA-approved labeling for additional information.		